

On Oscillation of Solutions of Higher Order Forced Functional Differential Inequalities

S. R. GRACE AND B. S. LALLI*

*Department of Mathematics, University of Saskatchewan,
Saskatoon, Saskatchewan S7N 0W0, Canada*

Submitted by W. Welsh

Oscillation criteria for the class of forced functional differential inequalities $x(t)\{L_n x(t) + f(t, x(t), x[g_1(t)], \dots, x[g_m(t)]) - h(t)\} \leq 0$, for n even, and $x(t)\{L_n x(t) - f(t, x(t), x[g_1(t)], \dots, x[g_m(t)]) - h(t)\} \geq 0$, for n odd, are established.

1. INTRODUCTION

A great deal of literature exists on the oscillation and nonoscillation of the homogeneous equation

$$x^{(n)}(t) + H(t, x[g(t)]) = 0$$

and the nonhomogeneous equation

$$x^{(n)} + H(t, x[g(t)]) = h(t),$$

both for n even and n odd. For this see [2-8] and the references cited in them. However, not much is known about inequalities of the type

$$x(t)\{L_n x(t) + f(t, x(t), x[g_1(t)], \dots, x[g_m(t)]) - h(t)\} \leq 0 \quad \text{for } n \text{ even} \quad (1)$$

and

$$x(t)\{L_n x(t) - f(t, x(t), x[g_1(t)], \dots, x[g_m(t)]) - h(t)\} \geq 0 \quad \text{for } n \text{ odd}, \quad (2)$$

where $L_0 x(t) = x(t)$, $L_k x(t) = a_k(t)(L_{k-1} x(t))'$, $a_n(t) = 1$, $k = 1, 2, \dots, n$, ($' = d/dt$).

Our main aim in this paper is to discuss the oscillatory behavior of the solutions of (1) (or (2)). We impose conditions on a_i , $i = 1, 2, \dots, n-1$ so

*The research of this author was supported partially by National Science Education and Research Council Grant A5293.

that we can replace the often used statement "the bounded solution $x(t)$ of (1) (or (2))" by "the solution $x(t)$ of (1) (or (2)) such that

$$\lim_{t \rightarrow \infty} \frac{x(t)}{a_2(t)} = 0, \quad a_2(t) = \int_c^t \frac{1}{a_1(s_1)} \int_c^{s_1} \frac{1}{a_2(s_2)} ds_2 ds_1,$$

and obtain new criteria for oscillation of solutions of (1) (or (2)).

Our results in Section 2 generalize and improve some of the results of the present authors [1], Kim [8], Lovelady [9], and Kartsatos and Manougian [7].

2. MAIN RESULTS

In the sequel it is assumed that

$$a_k, g_i \in C[R_+ = [0, \infty), R_+ \setminus \{0\}], \quad g_i(t) \rightarrow \infty \quad \text{as } t \rightarrow \infty, \quad (3)$$

for $k = 1, 2, \dots, n-1$ and $i = 1, 2, \dots, m$;

$$\int_c^\infty \frac{1}{a_k(s)} ds = \infty, \quad k = 1, 2, \dots, n-1; \quad (4)$$

$$\lim_{t \rightarrow \infty} \frac{1}{a_2(t)} \sum_{j=1}^k c_j a_j(t) > 0, \quad \alpha_0(t) = 1, \quad (5)$$

for every choice of constants c_j with $c_k > 0$, $k = 2, 3, \dots, n-1$, where

$$\begin{aligned} a_1(t) &= \int_c^t \frac{1}{a_1(s)} ds, \quad a_k(t) \\ &= \int_c^t \frac{1}{a_1(s_1)} \int_c^{s_1} \frac{1}{a_2(s_2)} \cdots \int_c^{s_{k-1}} \frac{1}{a_k(s_k)} ds_k ds_{k-1} \cdots ds_1, \end{aligned}$$

$k = 1, 2, \dots, n-1$ and $c \geq 0$;

$$f \in C[R_+ \times R^{m+1}, R] \quad \text{for } x, x_i > 0, \quad i = 1, 2, \dots, m \quad (6)$$

implies $f(t, x, x_1, \dots, x_m)$ is positive and nondecreasing with respect to x, x_i , $i = 1, 2, \dots, m$ for all $t \geq 0$;

$$f(t, x, x_1, \dots, x_m) \leq -f(t, -x, -x_1, \dots, -x_m), \quad (7)$$

for $x, x_i > 0$, $i = 1, 2, \dots, m$ and all $t \geq 0$;

$$h \in C[R_+, R] \quad (8)$$

and there exists an oscillatory function $\eta(t)$ such that $L_n \eta(t) = h(t)$, $L_k \eta(t)$ is oscillatory and $L_k \eta(t) \rightarrow 0$ as $t \rightarrow \infty$ for $k = 0, 1, \dots, n-1$.

The following Lemma is analogous to [1, Theorem 1]. We shall sketch the proof.

LEMMA. *Let conditions (3)–(8) hold. If x is a nontrivial solution of (1) (or (2)) such that*

$$x(t) > 0, \quad x(t)/\alpha_2(t) \rightarrow 0 \quad \text{as } t \rightarrow \infty, \quad t \geq t_0 \geq 0,$$

then there is a $t^ \geq t_0$ such that $y(t) = x(t) - \eta(t)$ is a solution of (1) (or (2)) for $t \geq t^*$ with the following properties:*

- (i) *we have $y(t) > 0$, $y(t)/\alpha_2(t) \rightarrow 0$ as $t \rightarrow \infty$;*
- (ii) *we have $(-)^{k-1} L_k y(t) > 0$, $L_k x(t) \cdot L_k y(t) > 0$ for $t \geq t^*$, $k = 1, 2, \dots, n-1$;*
- (iii) *we have $L_k x(t) \rightarrow 0$ monotonically as $t \rightarrow \infty$, $k = 2, 3, \dots, n-1$.*

Proof. We only consider (1). Let $x(t)$ be a nontrivial solution of (1) such that $x(t) > 0$ for $t \geq t_0 \geq 0$ and $x(t)/\alpha_2(t) \rightarrow 0$ as $t \rightarrow \infty$. Choose $t_1 \geq t_0$ so that $g_i(t) \geq t_0$, $t \geq t_1$, $i = 1, 2, \dots, m$. Thus $x[g_i(t)] > 0$ for $t \geq t_1$ and $i = 1, 2, \dots, m$. Inequality (1) becomes

$$L_n x(t) + f(t, x(t), x[g_1(t)], \dots, x[g_m(t)]) - h(t) \leq 0. \quad (9)$$

Let $y(t) = x(t) - \eta(t)$, then (9) becomes

$$L_n y(t) + f(t, y(t) + \eta(t), y[g_1(t)] + \eta[g_1(t)], \dots, y[g_m(t)] + \eta[g_m(t)]) \leq 0. \quad (10)$$

It is easy to check, using Kiguradze's lemma, that $L_k y(t)$ is of fixed sign for $t \geq t_1$ and $k = 0, 1, \dots, n$. Now if $y(t) < 0$ for $t \geq t_2 \geq t_1$, then $y(t) + \eta(t) > 0$ implies $\eta(t) > -y(t) > 0$, a contradiction to the oscillatory character of $\eta(t)$. Hence $y(t) > 0$ for $t \geq t_2$. We write (10) as the following system $y = y_1$, $y_1' = y_2/a_1, \dots, y_{n-1}' = y_n/a_{n-1}$ and $y_n'(t) \leq -f(t, y(t) + \eta(t), \dots, y[g_m(t)] + \eta[g_m(t)])$. As in the proof of [1, Theorem 1], one can easily see that $y_n(t) > 0$ for $t \geq t_3 \geq t_0$. If $L_{n-1} x(t) < 0$ for $t \geq t_3$, then $L_{n-1} y(t) + L_{n-1} \eta(t) < 0$ implies $-L_{n-1} \eta(t) > L_{n-1} y(t) > 0$, a contradiction to the oscillatory character of $L_{n-1} \eta(t)$. It is now easy to conclude that $y_n(t) \rightarrow 0$ as

$t \rightarrow \infty$ for $n > 2$. If this were not the case, there would exist a constant $C > 0$ such that

$$y_n(t) > C \quad \text{for } t \geq t_4, \quad \text{for some } t_4 \geq t_0.$$

However this implies that

$$\begin{aligned} y(t) = y_1(t) &> y_1(t_4) + y_2(t_4) \int_{t_4}^t \frac{1}{a_1(s)} ds + \dots \\ &+ y_{n-1}(t_4) \int_{t_4}^t \frac{1}{a_1(s_1)} \int_{t_4}^{s_1} \dots \\ &\times \int_{t_4}^{s_{n-3}} \frac{1}{a_{n-2}(s_{n-2})} ds_{n-2} \dots ds_1 \\ &+ C \int_{t_4}^t \frac{1}{a_1(s_1)} \int_{t_4}^{s_1} \dots \\ &\times \int_{t_4}^{s_{n-2}} \frac{1}{a_{n-1}(s_{n-1})} ds_{n-1} \dots ds_1 \\ &= \sum_{i=0}^{n-2} y_{i+1}(t_4) \alpha_i(t) + C \alpha_{n-1}(t), \quad \alpha_0(t) = 1. \end{aligned}$$

Dividing the above inequality by $\alpha_2(t)$, taking the limit as $t \rightarrow \infty$ and using condition (5), we get a contradiction to the fact that $y_1(t)/\alpha_2(t) \rightarrow 0$ as $t \rightarrow \infty$. Continuing this process we deduce that $y_n(t) > 0$, $y_{n-1}(t) < 0, \dots, y_2(t) > 0$, $y_1(t) > 0$ for $t \geq t_0$ and $y_k(t) \rightarrow 0$ as $t \rightarrow \infty$, $k = 3, 4, \dots, n$. This proves the Lemma.

In order to characterize the behaviors of solutions, we may reformulate the Lemma as follows:

THEOREM 1. *If x is a nontrivial solution of (1) (or (2)) such that $\lim_{t \rightarrow \infty} (x(t)/\alpha_2(t)) = 0$, then either*

- (a) *x is an oscillatory solution of (1) (or (2)), or else*
- (b) *$x \geq 0$ (≤ 0) on $[t_1, \infty)$ for some $t_1 \geq t_0$ and x ($-x$) satisfies the conclusion of the Lemma. In particular x ($-x$) increases (decreases) monotonically on $[t_1, \infty)$.*

COROLLARY. *If x is a nontrivial solution of (1) (or (2)) such that $x(t) \rightarrow 0$ as $t \rightarrow \infty$, then x is oscillatory.*

Remarks. (1) If $h(t) = 0$, $a_i(t) = 1$, $i = 1, \dots, n-1$, $f(t, x, \dots) = q(t)x$.

and the inequality in (1) (or (2)) be replaced by equality, then [8, Theorem 1] and our Lemma are the same.

(2) If $h(t) = 0$ and $f(t, x, \dots) = q(t)f(x[g(t)])$, then [1, Theorem 1] is included in our Lemma.

Let $1 \leq k \leq n-1$ and $t \in [t_0, \infty)$. We define

$$w_1(t) = \int_{t_0}^t \frac{1}{a_1(u)} du \quad \text{and} \quad w_k(t) = \int_{t_0}^t \frac{1}{a_k(u)} w_{k-1}(u) du.$$

THEOREM 2. *Let conditions (3), (4), and (6)–(8) hold, and let*

$$\int_{t_0}^{\infty} w_{n-1}(s) f(s, c, \dots, c) ds = \infty, \quad (11)$$

where c is a positive constant; then every bounded solution of (1) (or (2)) is oscillatory.

Proof. We only consider (1). Let $x(t)$ be a nonoscillatory bounded solution of (1). Without loss of generality, we may assume that $x(t)$ and $x[g_i(t)]$ are positive for $t \geq t_0$ and $i = 1, 2, \dots, m$. Hence

$$L_n x(t) + f(t, x(t), x[g_1(t)], \dots, x[g_m(t)]) - h(t) \leq 0.$$

Let $y(t) = x(t) - \eta(t)$. Then

$$\begin{aligned} &L_n y(t) + f(t, y(t) + \eta(t), y[g_1(t)] \\ &\quad + \eta[g_1(t)], \dots, y[g_m(t)] + \eta[g_m(t)]) \leq 0, \end{aligned}$$

which implies $L_n y(t) < 0$. By our Lemma, we have

$$\sum_{j=1}^{n-1} (-)^{j-1} w_j(t) L_j y(t) \geq 0, \quad t \geq t_1 \geq t_0.$$

It is easily verified that

$$\begin{aligned} y(t) &\geq y(t_1) + \sum_{j=1}^{n-1} (-)^{j-1} w_j(t) L_j y(t) - \sum_{j=1}^{n-1} (-)^{j-1} w_j(t_1) L_j y(t_1) \\ &\quad + \int_{t_1}^t w_{n-1}(s) f(s, y(s) + \eta(s), \dots, y[g_m(s)] + \eta[g_m(s)]) ds. \end{aligned}$$

Thus for $t \geq t_1$, we obtain

$$y(t) \geq y(t_1) - \sum_{j=1}^{n-1} (-)^{j-1} w_j(t_1) L_j y(t_1) + \int_{t_1}^t w_{n-1}(s) f(s, y(s)) \\ + \eta(s), \dots, y[g_m(s)] + \eta[g_m(s)] ds. \quad (12)$$

Let $g_*(t) = \min\{g_1(t), g_2(t), \dots, g_m(t)\}$. Since $x'(t) > 0$ and $y'(t) > 0$ for $t \geq t_1$, we have

$$x[g_i(t)] \geq x[g_*(t)] = y[g_*(t)] + \eta[g_*(t)] \\ \geq x[g_*(t_1)] = y[g_*(t_1)] + \eta[g_*(t_1)] \\ = c > 0 \quad \text{for } t \geq t_1,$$

and $i = 1, 2, \dots, m$. Therefore (12) becomes

$$y(t) \geq y(t_1) - \sum_{j=1}^{n-1} (-)^{j-1} w_j(t_1) L_j y(t_1) + \int_{t_1}^t w_{n-1}(s) f(s, c, \dots, c) ds.$$

Thus by (11), $\lim_{t \rightarrow \infty} y(t) = \infty$, a contradiction. This contradiction establishes our theorem.

THEOREM 3. *Let conditions (3)–(8) hold, and let*

$$\limsup_{t \rightarrow \infty} \frac{1}{\alpha_2(t)} \int_{t_1}^t w_{n-1}(s) f(s, c, \dots, c) ds > 0, \quad (13)$$

where c is a positive constant. Then every nontrivial solution $x(t)$ of (1) (or (2)) such that $x(t)/\alpha_2(t) \rightarrow 0$ as $t \rightarrow \infty$ is oscillatory.

Proof. The proof of Theorem 3 is similar to that of Theorem 2 and is omitted.

Remarks. (1) The above results hold equally well for the case when $h(t) = 0$.

(2) If $h(t) = 0$, then [9, Theorem 1] and [1, Theorem 4] are included, respectively, in our Theorems 2 and 3.

EXAMPLE. The equation

$$\left(\frac{1}{t} \left(\frac{1}{t} \left(\frac{1}{t} x' \right)' \right)' \right) + \frac{231}{16t^7} x^\alpha(t^{1/\alpha}) = 0, \quad t > 0, \quad (14)$$

where α is the quotient of two odd positive integers, has the nonoscillatory unbounded solution $x(t) = \sqrt{t}$. All the conditions of Theorem 2 are satisfied,

thus all bounded solutions of (14) are oscillatory. Only condition (13) of Theorem 3 is violated. We may add that the solutions of the equation

$$\left(\frac{1}{t} \left(\frac{1}{t} \left(\frac{1}{t} x' \right)' \right)' \right) + \frac{231}{16t^7} x^\alpha (t^{1/\alpha}) = e^{-t} \sin t, \quad t > 0. \quad (15)$$

α as above, have the same oscillation property as those of (14) for t sufficiently large because h satisfies condition (8).

For convenience of notation for any $t_0 \geq 0$ and $t \geq t_0$ we let

$$w(t) = \max_{1 \leq i \leq m} \alpha_i(t), \quad \gamma(t) = \int_{t_0}^t \frac{1}{a_1(s_1)} \int_{t_0}^{s_1} \cdots \int_{t_0}^{s_{n-1}} h(s) ds ds_{n-1} \cdots ds_1.$$

THEOREM 4. *Let conditions (3) and (6) hold, and let*

$$\limsup_{t \rightarrow \infty} [\gamma(t) + kw(t)] = +\infty,$$

$$\liminf_{t \rightarrow \infty} [\gamma(t) + kw(t)] = -\infty.$$

for $t \in [0, \infty)$ and every $k \in (0, \infty)$. Then every solution of (1) (or (2)) is oscillatory.

Proof. The proof is similar to that of [7, Theorem 2.1] and is omitted.

THEOREM 5. *Let conditions (3), (4), and (6) hold, and*

$$\liminf_{t \rightarrow \infty} \int_0^t w_{n-1}(s) [-f(s, k, k, \dots, k) + h(s)] ds = -\infty, \quad (16)$$

$$\limsup_{t \rightarrow \infty} \int_0^t w_{n-1}(s) [-f(s, -k, \dots, -k) + h(s)] ds = \infty, \quad (17)$$

for every $k > 0$. Moreover, let

$$\int h(t) dt = \phi(t) + C, \quad \int \frac{\phi(t)}{a_{n-1}(t)} dt = \phi_1(t) + C, \quad (18)$$

where C is any arbitrary constant, ϕ and ϕ_1 are two bounded functions with ϕ oscillatory.

Then any bounded solution of (1) (or (2)) is oscillatory.

Proof. The proof is an adaptation of an argument developed by Kartsatos and Manougian [7, Theorem 2.4]. Again we only consider (1). Let

$x(t)$ be a nonoscillatory solution of (1). Without loss of generality, we assume that $x(t)$, $x[g_i(t)]$, $i = 1, 2, \dots, m$ are positive, $t \geq t_1 \geq t_0$. We also assume that there exists $L > 0$ such that $0 < x(t) \leq L$ and $0 < x[g_i(t)] \leq L$ for $t \geq t_1$, $i = 1, 2, \dots, m$. We may take t_2 such that $\phi(t_2) = 0$. Now consider the transformation

$$M(t) = L_{n-1}x(t) - \phi(t), \quad t \geq t_2.$$

Then we have

$$M'(t) = -f(t, x(t), \dots, x[g_m(t)]) \leq 0.$$

Thus $M(t)$ is decreasing in $[t_2, \infty)$. Assume that $M(t_3) = -\lambda < 0$ for some $t_3 \geq t_2$. Then $M(t) \leq -\lambda$ in $[t_3, \infty)$. Thus

$$\frac{-\lambda}{a_{n-1}(t)} \geq \frac{M(t)}{a_{n-1}(t)} = (L_{n-2}x(t))' - \frac{\phi(t)}{a_{n-1}(t)}.$$

Thus

$$-\lambda \int_{t_3}^t \frac{1}{a_{n-1}(s)} ds \geq L_{n-2}x(t) - L_{n-2}x(t_3) - \phi_1(t) + \phi_1(t_3).$$

Since $\phi_1(t)$ is bounded, $\lim_{t \rightarrow \infty} L_{n-2}x(t) = -\infty$, a contradiction to the positivity of $x(t)$. Thus $M(t) \geq 0$ for $t \in [t_2, \infty)$. Now, by integration of the function

$$-(L_n x(t) - \phi'(t)) = f(t, x(t), x[g_1(t)], \dots, x[g_m(t)])$$

we obtain

$$-M(t) + L_{n-1}x(t_2) = \int_{t_2}^t f(s, x(s), \dots, x[g_m(s)]) ds,$$

from which

$$L_{n-1}x(t_2) \geq \int_{t_2}^t f(s, x(s), \dots, x[g_m(s)]) ds.$$

Since t_2 is arbitrary, we have $L_{n-1}x(t) \geq 0$ for any $t \geq t_2$. It follows from our Lemma that

$$(-)^k L_k x(t) \leq 0 \quad \text{for } t \geq \bar{t} \geq t_2, \quad k = 1, \dots, n-1. \quad (19)$$

This implies that $x'(t) \geq 0$ for $t \geq \bar{t}$. Thus there exists $\bar{t}_1 \geq \bar{t}$ such that $x[g_i(t)] \geq K > 0$, $x(t) \geq K > 0$ for every $t \geq \bar{t}_1$, where K is a constant. Now

differentiating the function $T(t) = w_{n-1}(t) L_{n-1}x(t)$, $t \geq \bar{t}_1$, and integrating from \bar{t}_1 to $t \geq \bar{t}_1$, we obtain

$$\begin{aligned} T(t) &= T(\bar{t}_1) + \int_{\bar{t}_1}^t w_{n-1}(s) [-f(s, x(s), \dots, x(g_m(s))) + h(s)] ds \\ &\quad + \int_{\bar{t}_1}^t \frac{w_{n-2}(s)}{a_{n-1}(s)} L_{n-1}x(s) ds \\ &\leq T(\bar{t}_1) + \int_{\bar{t}_1}^t w_{n-1}(s) [-f(s, K, \dots, K) + h(s)] ds \\ &\quad + \int_{\bar{t}_1}^t w_{n-2}(s) (L_{n-2}x(s))' ds. \end{aligned}$$

Now, applying (16), there exists a sequence $\{t_l\}$, $l = 1, 2, \dots$ such that $\lim_{l \rightarrow \infty} t_l = \infty$ and

$$\lim_{l \rightarrow \infty} \int_{t_l}^{t_l'} w_{n-1}(s) [-f(s, K, K, \dots, K) + h(s)] ds = \infty.$$

This implies that

$$\int_{\bar{t}_1}^{\infty} w_{n-2}(s) (L_{n-2}x(s))' ds = \infty \quad (20)$$

because $L_{n-1}x(t) \geq 0$, $t \in [\bar{t}_1, \infty)$. Successive integration by parts of (20) and by the use of (19) we obtain

$$\begin{aligned} \int_{\bar{t}_1}^{\infty} w_{n-i}(s) (L_{n-i}x(s))' ds &= +\infty, & \text{if } i = 2, 4, \dots, n. \\ &= -\infty, & \text{if } i = 3, 4, \dots, n-1. \end{aligned}$$

Thus if $i = n$, $\lim_{t \rightarrow \infty} x(t) = \infty$, a contradiction to the fact that x is a bounded function.

THEOREM 6. *Let conditions (3)–(6) and (18) hold, and*

$$\liminf_{t \rightarrow +\infty} \frac{1}{\alpha_2(t)} \int_0^t w_{n-1}(s) [-f(s, k, \dots, k) + h(s)] ds < 0, \quad (21)$$

$$\limsup_{t \rightarrow +\infty} \frac{1}{\alpha_2(t)} \int_0^t w_{n-1}(s) [-f(s, -k, \dots, -k) + H(s)] ds > 0, \quad (22)$$

for every $k > 0$. Then every nontrivial solution $x(t)$ of (1) (or (2)) such that $x(t)/\alpha_2(t) \rightarrow 0$ as $t \rightarrow \infty$ is oscillatory.

Proof. The proof is similar to the proof of Theorem 5 and is omitted.

THEOREM 7. *Let conditions (3), (4), (6), and (18) hold, and*

$$\liminf_{t \rightarrow \infty} \int_0^t [-f(s, k, \dots, k) + h(s)] ds = -\infty,$$

$$\limsup_{t \rightarrow \infty} \int_0^t [-f(s, -k, \dots, -k) + h(s)] ds = +\infty.$$

Then every solution of (1) (or (2)) is oscillatory.

Proof. Let $x(t)$ be a nonoscillatory solution of (1). Without loss of generality we may assume that $x(t)$, $x[g_i(t)]$, $i = 1, \dots, m$ are positive for $t \geq t_1 \geq t_0$. Thus it follows from the proof of Theorem 5 that $L_{n-1}x(t) \geq 0$ for every $t \geq t_2$, for some $t_2 \geq t_1$. This implies that there exists a constant $K > 0$ such that $x(t) \geq K$, $x[g_i(t)] \geq K$ for every $t \geq t_2$. This follows from the fact that $x(t)$ is positive, n is even, and $L_{n-1}x(t) \geq 0$, $x'(t) \geq 0$ for all large t . Thus integrating (1) once we get

$$\begin{aligned} L_{n-1}x(t) &= L_{n-1}x(t_2) + \int_{t_2}^t [-f(s, x(s), \dots, x[g_m(s)]) + h(s)] ds \\ &\leq L_{n-1}x(t_2) + \int_{t_2}^t [-f(s, K, \dots, K) + h(s)] ds, \end{aligned}$$

which implies that $\liminf_{t \rightarrow \infty} x(t) = -\infty$, a contradiction. A similar argument holds for $x(t)$ eventually negative and this completes the proof.

Remarks. (1) If $a_i(t) = 1$, $i = 1, 2, \dots, n-1$, then [7, Theorems 2.4 and 2.6] are included, respectively, in our Theorems 6 and 7.

(2) We believe that our Theorems 3 and 6 are new oscillation criteria for (1) (or (2)).

REFERENCES

1. S. R. GRACE AND B. S. LALLI, On oscillation of solutions of n th order delay differential equation, *J. Math. Anal. Appl.* **91** (1983), 328-339.
2. W. E. MAHFOUD, Characterization of oscillation of solutions of the delay equation $x^{(n)} + a(t)f(x[q(t)]) = 0$, *J. Differential Equations* **28** (1978), 437-451.
3. W. E. MAHFOUD, Oscillation and asymptotic behavior of solutions of n th order nonlinear delay differential equation, *J. Differential Equations* **24** (1977), 617-627.
4. A. G. KARTSATOS, On the maintenance of oscillations under the effect of a small forcing term, *J. Differential Equations* **10** (1970), 355-363.
5. A. G. KARTSATOS, Maintenance of oscillations under the effect of a periodic forcing term, *Proc. Amer. Math. Soc.* **33** (1972), 377-383.

6. A. G. KARTSATOS, Recent results on oscillation of solutions of forced and perturbed differential equations of even order in "Proceedings, NSF-CBNS Reg. Conference, Stability of Dynamical Systems, Mississippi State University, 1977" (J. R. Graef, Ed.), Dekker, New York, 1977.
7. A. G. KARTSATOS AND M. N. MANOUGIAN, Further results on oscillation of functional differential equation. *J. Math. Anal. Appl.* **53** (1976), 28-37.
8. W. J. KIM, Monotone and oscillatory solutions of $y(n) + py = 0$. *Proc. Amer. Math. Soc.* **62** (1977), 77-82.
9. D. L. LOVELADY, On oscillatory behavior of bounded solutions of higher order differential equations. *J. Differential Equations* **19** (1975), 167-175.